

Models for jet power in elliptical galaxies: support for rapidly spinning black holes

R. S. Nemmen¹, R. G. Bower², A. Babul³, and T. Storchi-Bergmann¹

¹*Instituto de Física, Universidade Federal do Rio Grande do Sul (Brazil)*

²*Department of Physics, Institute for Computational Cosmology, Durham University (UK)*

³*Department of Physics & Astronomy, University of Victoria (Canada)*

Abstract. Recently, Allen et al. measured a tight correlation between the Bondi accretion rates and jet powers of the nuclei of nearby X-ray luminous elliptical galaxies. We employ two models of jet powering to understand the above correlation and derive constraints on the spin and accretion rate of the central black holes. The first is the Blandford-Znajek model, in which the spin energy of the hole is extracted as jet power; the second model is an hybrid version of the Blandford-Payne and Blandford-Znajek processes, in which the outflow is generated in the inner parts of the accretion disk. We assume advection-dominated accretion flows (ADAF) and account for general relativistic effects. Our modelling implies that for typical values of the disk viscosity parameter $\alpha \sim 0.01 - 1$ the tight correlation implies the narrow range of spins $j \approx 0.7 - 1$ and accretion rates $\dot{M}(R_{\text{ms}}) \approx (0.01 - 1)\dot{M}_{\text{Bondi}}$. Our results provide support for the “spin paradigm” scenario and suggest that the central black holes in the cores of clusters of galaxies must be rapidly rotating in order to drive radio jets powerful enough to quench the cooling flows.

1. Introduction

Recently, Allen et al. (2006) (hereafter A06) measured a tight correlation between the Bondi accretion rates $\dot{M}_{\text{Bondi}}$ and jet powers P_{jet} of the nuclei of nine nearby, X-ray luminous elliptical galaxies using *Chandra* X-ray observations. It is imperative to understand how this correlation is established and what constraints on the properties of the central black holes can be derived from it. This is the goal of the present work.

2. Models for the jet power: constraints on the black hole spin and accretion rate

The origin of jets is likely to be related to the spin of the central black hole (e.g., Blandford 2002). With this in mind, we employ two physical models of jet production which relate the spin j ($\equiv a/M_{\bullet}$) and accretion rate $\dot{M}$ onto the black hole to the observed jet power in order to understand the empirical correlation of A06: the Blandford-Znajek (BZ) model (e.g., Blandford 2002), in which magnetic fields threading the hole extract its rotational energy and drive the jet; and a “hybrid model” (Meier 2001) which combines the Blandford-Payne




Figure 1. *Left*: comparison of the spin dependence of P_{jet} predicted by the models for two values of α , taking $\dot{M}_{\text{ms}} = \dot{M}(R_{\text{ms}}) = 1.6 \times 10^{-3} M_{\odot} \text{ yr}^{-1}$. The right axis shows the jet efficiency $\eta_{\text{jet}} \equiv P_{\text{jet}}/\dot{M}_{\text{ms}}c^2$. *Right*: contours of the model parameters $\epsilon_{\text{Bondi}} = \dot{M}_{\text{ms}}/\dot{M}_{\text{Bondi}}$ and j which reproduce the measured values of A ($A = 0.69 \pm 0.19$, labels beside each contour) of the correlation of A06, where $\log(P_{\text{Bondi}}/10^{43} \text{ erg s}^{-1}) = A + B \log(P_{\text{jet}}/10^{43} \text{ erg s}^{-1})$ ($B = 1$ from the jet models).

and BZ mechanisms, in which the fields tap energy from the accretion flow and the spinning hole. We assume that in the nuclei of the observed elliptical galaxies the accretion flow around the black hole is advection-dominated (advection-dominated accretion flows, hereafter ADAF, e.g., Nemmen et al. 2006, see also contribution of F. Yuan in this volume). ADAFs are expected to be associated with the production of radio jets (e.g., Meier 2001). We take into account general relativistic effects not fully appreciated before in such models, in particular the dependence of the radius of the marginally stable orbit (R_{ms}) on the black hole spin, and frame-dragging effects that intensify the magnetic field strength. Figure 1 (left) shows the spin dependence of P_{jet} we obtain from the modelling.

Figure 1 (right) illustrates the accretion rates and spins required to reproduce the correlation between $\dot{M}_{\text{Bondi}}$ and P_{jet} for $\alpha = 0.3$. For typical values of the viscosity parameter $\alpha \sim 0.01 - 1$, we obtain that the tight correlation of A06 implies the narrow range of spins $j \approx 0.7 - 1$ and accretion rates $\dot{M}_{\text{ms}} \approx (0.01 - 1)\dot{M}_{\text{Bondi}}$. These results are discussed in more depth in Nemmen et al. 2007.

References

- Allen, S. W., et al. 2006, MNRAS, 372, 21
- Blandford, R. D. 2002, in Lighthouses of the Universe, ed. M. Gilfanov, R. Sunyaev, & E. Churazov (Berlin: Springer), 381
- Meier, D. L. 2001, ApJ, 548, L9
- Nemmen, R. S., et al. 2006, ApJ, 643, 652
- Nemmen, R. S., Bower, R. G., Babul, A., & Storchi-Bergmann, T. 2007, MNRAS, submitted (astro-ph/0612354)